

**LE/AF Purified Anti-Mouse CD8a (53.6.7)
Monoclonal Antibody**天津三箭生物技术股份有限公司
Tianjin Sungene Biotech Co., Ltd.
标准 高效 稳定 Precision Efficient Stable

Catalog Number	Vial Size
M10081-14B	50 µg
M10081-14E	500 µg
M10081-14F	1 mg

Market | 400-621-0003
marketing@sungenebiotech.com**Support** | 022-66211636-8024
techsupport@sungenebiotech.com**Web** | www.sungenebiotech.com

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.
This product is guaranteed up to one year from purchase.

Purified Antibody Characterization

Clone	Isotype	Reactivity
53.6.7	Rat IgG2a	Mouse

Description

CD8, also known as Lyt-2, Ly-2, or T8, consists of disulfide-linked α and β chains that form the α (CD8a)/ β (CD8b) heterodimer and γ/δ homodimer. CD8a is a 34 kD protein that belongs to the immunoglobulin family. The CD8 α/β heterodimer is expressed on the surface of most thymocytes and a subset of mature TCR α/β T cells. CD8 expression on mature T cells is non-overlapping with CD4. The CD8 α/α homodimer is expressed on a subset of γ/δ TCR-bearing T cells, NK cells, intestinal intraepithelial lymphocytes, and lymphoid dendritic cells. CD8 is an antigen co-receptor on T cells that interacts with MHC class I on antigen-presenting cells or epithelial cells. CD8 promotes T cell activation through its association with the TCR complex and protein tyrosine kinase Lck.

Reported Applications

This 53.6.7 antibody has been reported for use in Flow Cytometric Analysis, Immunohistochemical Staining of Frozen Tissue Sections, Immunoprecipitation. It has also been reported for use in cell depletion.

Product Information**Production Method:** Stirred tank fermentation**Medium:** Hybridoma-SFM + 1%FCS + Glc + P/S**Purification Method:** Protein G**Concentration:** 1 mg/ml**Endotoxin:** < 2.00 EU/mg (LAL)**Purity:** >95% (by SDS-PAGE)**Sterile:** 0.2 µm Filtration**Formulated:** PBS, pH7.2**Storage:** Keep as concentrated solution. Store at 4°C as an undiluted liquid. For extended storage aliquot contents and freeze at -20°C or lower. Avoid cycles of freezing and thawing.**For Research Use Only.**