PE/Cy5 Anti-Rat CD4 (domain 1) Monoclonal Antibody

Vial Size

25 µg

100 µg

| | 天津三箭生物技术股份有限公司 Tianjin Sungene Biotech Co., Ltd. ^{精准 高效 稳定} Precision Efficient Stable |
|---------|---|
| Market | 400-621-0003 marketing@sungenebiotech.com |
| Support | 022-66211636-8024 techsupport@sungenebiotech.com |
| Web | www.sungenebiotech.com |

Important Note: Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Purified Antibody Characterization

R10041-35A

R10041-35C

| Clone | Isotype | Reactivity | |
|-------|-------------|------------|--|
| OX-38 | Mouse IgG2a | Rat | |

Description

CD4, also known as T4, is a 55kD glycoprotein member of the immunoglobin superfamily and is expressed on majority of thymocytes, macrophages, and a peripheral T cell subset (T helper cells). CD4 is a T cell co-receptor that interacts with the MHC class II molecule and is involved in T cell activation.

Illustration of Immunofluorescent Staining



FITC anti-rat CD8

Wister rat splenocytes stained with PE/ Cy5 anti-rat CD4 and FITC anti-rat CD8 antibody

Product Information

Conjugation: PE/Cy5

Formulation: PBS pH 7.2, 0.09% NaN₃, 0.2% BSA

Concentration: 0.2 mg/ml

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged exposure to light. **Do not freeze.**

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used ≤ 0.25 µg /10⁶ cells in 100 µl). Since applications vary, the appropriate dilutions must be determined for individual use.

References

[1] Pettigrew, G.J., et al. 1998. J. Immunol. 161:1292.

For Research Use Only.